

## Week 7 Chapter 4 and 6

4-76. a)  $P(X \leq 0) = \int_0^0 \lambda e^{-\lambda x} dx = 0$

b)  $P(X \geq 2) = \int_2^{\infty} 2e^{-2x} dx = -e^{-2x} \Big|_2^{\infty} = e^{-4} = 0.0183$

c)  $P(X \leq 1) = \int_0^1 2e^{-2x} dx = -e^{-2x} \Big|_0^1 = 1 - e^{-2} = 0.8647$

d)  $P(1 < X < 2) = \int_1^2 2e^{-2x} dx = -e^{-2x} \Big|_1^2 = e^{-2} - e^{-4} = 0.1170$

e)  $P(X \leq x) = \int_0^x 2e^{-2t} dt = -e^{-2t} \Big|_0^x = 1 - e^{-2x} = 0.05$  and  $x = 0.0256$

4-84. Let  $X$  denote the time until a message is received. Then,  $X$  is an exponential random variable and  $\lambda = 1/E(X) = 1/2$ .

a)  $P(X > 2) = \int_2^{\infty} \frac{1}{2} e^{-x/2} dx = -e^{-x/2} \Big|_2^{\infty} = e^{-1} = 0.3679$

b) The same as part a.

c)  $E(X) = 2$  hours.

4-90. Let  $Y$  denote the number of arrivals in one hour. If the time between arrivals is exponential, then the count of arrivals is a Poisson random variable and  $\lambda = 1$  arrival per hour.

a)  $P(Y > 3) = 1 - P(Y \leq 3) = 1 - \left[ \frac{e^{-1}1^0}{0!} + \frac{e^{-1}1^1}{1!} + \frac{e^{-1}1^2}{2!} + \frac{e^{-1}1^3}{3!} \right] = 0.01899$

b) From part a),  $P(Y > 3) = 0.01899$ . Let  $W$  denote the number of one-hour intervals out of 30 that contain more than 3 arrivals. By the memoryless property of a Poisson process,  $W$  is a binomial random variable with  $n = 30$  and  $p = 0.01899$ .

$$P(W = 0) = \binom{30}{0} 0.01899^0 (1 - 0.01899)^{30} = 0.5626$$

c) Let  $X$  denote the time between arrivals. Then,  $X$  is an exponential random variable with

$$\lambda = 1 \text{ arrivals per hour. } P(X > x) = 0.1 \text{ and } P(X > x) = \int_x^{\infty} 1e^{-1t} dt = -e^{-1t} \Big|_x^{\infty} = e^{-1x} = 0.1.$$

Therefore,  $x = 2.3$  hours.

4-130 The time to failure (in hours) for a laser in a cytometry machine is modeled by an exponential distribution with 0.00004.

a)  $P(X > 20,000) = \int_{20000}^{\infty} 0.00004e^{-0.00004x} dx = -e^{-0.00004x} \Big|_{20000}^{\infty} = e^{-0.8} = 0.4493$


$$b) P(X < 30,000) = \int_{30000}^{\infty} 0.00004e^{-0.00004x} dx = -e^{-0.00004x} \Big|_0^{30000} = 1 - e^{-1.2} = 0.6988$$

c)

$$P(20,000 < X < 30,000) = \int_{20000}^{30000} 0.00004e^{-0.00004x} dx$$


$$= -e^{-0.00004x} \Big|_{20000}^{30000} = e^{-0.8} - e^{-1.2} = 0.1481$$

6-66.



It appears that the data do not come from a normal distribution. Very few of the data points fall on the line.

6-72.




The data appear to be normally distributed. Nearly all of the data points fall very close to, or on the line.

6-94. a) Descriptive Statistics

| Variable | N  | N* | Mean   | SE Mean | StDev  | Variance |
|----------|----|----|--------|---------|--------|----------|
| Density  | 29 | 0  | 5.4197 | 0.0629  | 0.3389 | 0.1148   |

| Variable | Minimum | Q1     | Median | Q3     | Maximum |
|----------|---------|--------|--------|--------|---------|
| Density  | 4.0700  | 5.2950 | 5.4600 | 5.6150 | 5.8600  |



b) There does appear to be a low outlier in the data.

c) Due to the very low data point at 4.07, the mean may be lower than it should be. Therefore, the median would be a better estimate of the density of the earth. The median is not affected by a few outliers.