

Math 341, Final Exam, Spring 2010, Name_____

Student # _____

Friday, May 14. *Please show complete solution (with all steps) to each problem to receive full credit.*

1. The Rockwell hardness index for steel is determined by pressing a diamond point into the steel and measuring the depth of penetration. For 50 specimens of an alloy of steel, the Rockwell hardness index average 61 with standard deviation 7. The manufacturer claims that this alloy has an average hardness index of at least 63. Is there sufficient evidence to refute the manufacturer's claim? Test at $\alpha = 0.01$ using a one-sided 99% confidence interval. Clearly specify the null and alternate hypotheses and the conclusion. (15 points)
2. The true mean manual dexterity scores of the second graders who participated in sports (μ_1) is being compared with that of second graders who did not participate in sports (μ_2). One wishes to establish that the true mean dexterity score, for second graders who participate in sports, is higher than that of those who did not participate in sports. If the sample sizes for two groups are taken to be the same and the sample variances are $s_1^2 = 4.3$ and $s_2^2 = 4.6$ (from pilot study), respectively. Find the sample sizes that give $\alpha = 0.01$ and $\beta = 0.15$, when $\mu_1 - \mu_2 = 2$. (10 points)

3. The hourly wages in a particular industry are normally distributed with mean \$13.20 and standard deviation \$3.00. A company in this industry employs 45 workers, paying them an average of \$11.50 per hour. (a) Can this company be accused of paying substandard wages? Use an $\alpha = 0.02$ level test. (b) Compute the type II error probability when $\mu = 13$. (18 points, 9 points each)

4. The geometric probability mass function is given by $p(y | p) = p(1 - p)^{y-1}$, $y = 1, 2, \dots$. A random sample of size n is taken from a population with a geometric distribution. Find the maximum likelihood estimator for p . (10 points)

5. Suppose that X_1, \dots, X_n and Y_1, \dots, Y_n are independent random samples from populations with means μ_1 and μ_2 and variances $0 < \sigma_1^2 < \infty$ and $0 < \sigma_2^2 < \infty$, respectively. Show that $\bar{X} - \bar{Y}$ is a consistent estimator of $\mu_1 - \mu_2$. (12 points)

6. Let Y_1, \dots, Y_n denote a random sample from the uniform distribution on the interval

$(\theta, \theta + 1)$. Let $\hat{\theta}_1 = \bar{Y} - \frac{1}{2}$ and $\hat{\theta}_2 = Y_{(n)} - \frac{n}{n+1}$.

a. Show that $\hat{\theta}_1$ and $\hat{\theta}_2$ are unbiased estimators of θ .

b. Find the efficiency of $\hat{\theta}_1$ relative to $\hat{\theta}_2$. (15 points)

7. Seasonal ranges (in hectors) for alligators were monitored on a lake outside Gainesville, Florida, by biologists from the Florida Game and Fish Commission. Four alligators monitored in the spring showed ranges of 8.0, 12.1, 8.1, and 18.2. Three different alligators monitored in the summer showed ranges of 102.0, 81.7, and 54.7. (a) Estimate the difference between mean spring and summer ranges, with a 99% confidence interval. (b) Are the two means different at level $\alpha = 0.01$? Why or why not? (c) What statistical assumptions did you make? (20 points)