

Numerical Analysis August 2019 Previous Qual

Question 6

José L. Pabón

Spring 2020

This work is based on the course textbook [1], the material discussed in lectures and office hours related to our course MAT614 and additional references. This is for the May 2018 qualifying exam.

1 Problem August 2019 3:

1.1 REDACTED

with respect to the weight function $w(x) = \frac{1}{\sqrt{1-x^2}}$.

The Chebyshev polynomials are of the form:

$$T_n(x) = \cos(n \arccos(x)).$$

We consider the weighted inner product for positive integers $m, n \in \mathbf{N}$:

$$\langle T_n(x), T_m(x) \rangle = \int_{-1}^1 T_n(x) T_m(x) w(x) dx.$$

We define $\theta = \arccos(x) \implies d\theta = -\frac{1}{\sqrt{1-x^2}} dx$, and substitute into our integral:

$$\int_{-1}^1 T_n(x) T_m(x) w(x) dx = - \int_{\pi}^0 \cos(n\theta) \cos(m\theta) d\theta = \int_0^{\pi} \cos(n\theta) \cos(m\theta) d\theta.$$

The integrand functions are an orthogonal family of functions which will equal to zero unless n equals to m .

$$\frac{1}{2} \int_0^{\pi} \cos((n+m)\theta) + \cos((n-m)\theta) d\theta = \frac{1}{2} \left[\frac{1}{(n+m)} \sin((n+m)\theta) + \frac{1}{(n-m)} \sin((n-m)\theta) \right] \Big|_0^{\pi}.$$

Thus we have that:

$$\frac{1}{2} \left[\frac{1}{(n+m)} \sin((n+m)\theta) + \frac{1}{(n-m)} \sin((n-m)\theta) \right] \Big|_0^\pi = \begin{cases} \pi, & \text{for } m = n = 0 \\ \frac{1}{2}\pi, & \text{for } m = n \neq 0 \\ 0, & \text{otherwise.} \end{cases}$$

1.2 REDACTED

Show that $T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x), n \geq 2$.

The left hand side is:

We again define $\theta = \arccos(x) \implies \cos \theta = x$

$$T_{n+1}(x) = \cos((n+1) \arccos(x)) \implies T_{n+1}(\theta) = \cos((n+1)\theta).$$

The right hand side we have is:

$$2xT_n(x) - T_{n-1}(x) \implies 2 \cos \theta \cos(n\theta) - \cos((n-1)\theta).$$

Then:

$$2 \cos \theta \cos(n\theta) = \cos(n+1)\theta - \cos(n-1)\theta.$$

Thus:

$$2 \cos \theta \cos(n\theta) - \cos((n-1)\theta) = \cos(n+1)\theta.$$

$$\therefore T_{n+1}(\theta) = \cos((n+1)\theta) = 2 \cos \theta \cos(n\theta) - \cos((n-1)\theta) \implies$$

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x) \checkmark.$$

1.3 REDACTED

Derive the three point Chebyshev Gauss quadrature formula:

$$I(f) = \int_{-1}^1 f(x) \frac{1}{\sqrt{1-x^2}} dx \approx \sum_{i=1}^3 w_i f(x_i).$$

We begin by computing the three roots of our $n = 3$ Chebyshev polynomial:

$$T_n(x) = \cos(n \arccos(x)) \text{ or } T_n(\theta) = \cos(n\theta).$$

Via the previous identities derived in the previous parts of this problem:

$$T_3(x) = \cos(3 \arccos(x)) \implies 2xT_2(x) - T_1(x) \implies 2x \cos(2 \arccos(x)) - \cos(1 \arccos(x)).$$

We have that $\cos(1 \arccos(x)) = x$. We apply the same recursion formula to our term $\cos(2 \arccos(x))$:

$$\cos(2 \arccos(x)) = 2x \cos(1 \arccos(x)) - \cos(0 \arccos(x)).$$

Thus, we have that:

$$T_3(x) = 2x(2x^2 - 1) - x = 4x^3 - 2x - x = 4x^3 - 3x.$$

This gives us our collocation points:

$$x_1 = -\frac{1}{2}\sqrt{3}, x_2 = 0, x_3 = \frac{1}{2}\sqrt{3}.$$

We form our system of equations with our monomial basis $1, x, x^2$:

For $f(x) = 1$ we have that:

$$\int_{-1}^1 \frac{1}{\sqrt{1-x^2}} dx \approx w_1 f(x_1) + w_2 f(x_2) + w_3 f(x_3) = w_1 + w_2 + w_3.$$

$$\int_{-1}^1 \frac{1}{\sqrt{1-x^2}} dx = \arcsin 1 - \arcsin -1 = \frac{1}{1}\pi - (-\frac{1}{1}\pi) = \pi$$

. Thus:

$$w_1 + w_2 + w_3 = \pi$$

For $f(x) = x$ we have that:

$$\int_{-1}^1 x \frac{1}{\sqrt{1-x^2}} dx \approx w_1 f(x_1) + w_2 f(x_2) + w_3 f(x_3) = w_1 - \frac{1}{2}\sqrt{3} + w_2 0 + w_3 \frac{1}{2}\sqrt{3}.$$

$$\int_{-1}^1 x \frac{1}{\sqrt{1-x^2}} dx = \arcsin 1 - \arcsin -1 = \text{An odd integrand over a symmetric interval} = 0$$

. Thus:

$$0 = w_1 - \frac{1}{2}\sqrt{3} + w_2 0 + w_3 \frac{1}{2}\sqrt{3}.$$

We multiply this equation by $2\sqrt{3}$ on both sides to find:

$$-3w_1 + 0w_2 + 3w_3 = 0.$$

For $f(x) = x^2$ we have that:

$$\int_{-1}^1 x^2 \frac{1}{\sqrt{1-x^2}} dx \approx w_1 f(x_1) + w_2 f(x_2) + w_3 f(x_3) = w_1 \left(-\frac{1}{2}\sqrt{3}\right)^2 + w_2 0 + w_3 \left(\frac{1}{2}\sqrt{3}\right)^2.$$

Integrating by parts we have that:

$$\int x^2 \frac{1}{\sqrt{1-x^2}} dx = \frac{1}{2}\pi.$$

Thus:

$$\frac{1}{2}\pi = w_1 \left(-\frac{1}{2}\sqrt{3}\right)^2 + w_2 0 + w_3 \left(\frac{1}{2}\sqrt{3}\right)^2 = w_1 \left(\frac{1}{4}3\right) + 0 + w_3 \left(\frac{1}{4}3\right)$$

Equivalently, we multiply this resulting equation by 4 on both sides, we have that:

$$2\pi = 3w_1 + 0 + 3w_3.$$

Thus we have our systems of equations:

$$w_1 + w_2 + w_3 = \pi$$

$$-3w_1 + 0 + 3w_3 = 0.$$

$$3w_1 + 0 + 3w_3 = 2\pi.$$

We solve this system of equations to find that our weights are:

$$w_1 = w_2 = w_3 = \frac{1}{3}\pi.$$

1.4 d - Approximate $I(f) = e^{-2x^2}$.

Thus, by our previous result, we know that:

$$I(f) \approx \sum_{i=1}^3 w_i f(x_i) = \frac{1}{3}\pi f(x_1) + \frac{1}{3}\pi f(x_2) + \frac{1}{3}\pi f(x_3) = \frac{1}{3}\pi e^{-2(-\sqrt{3}\frac{1}{2})^2} + \frac{1}{3}\pi e^{-2(0)^2} + \frac{1}{3}\pi e^{-2(\sqrt{3}\frac{1}{2})^2}.$$

Then we have that:

$$\frac{1}{3}\pi e^{-2(-\sqrt{3}\frac{1}{2})^2} + \frac{1}{3}\pi e^{-2(0)^2} + \frac{1}{3}\pi e^{-2(\sqrt{3}\frac{1}{2})^2} = \frac{1}{3}\pi(1 + e^{-6(\frac{1}{4})} + e^{-6(\frac{1}{4})}) = \frac{1}{3}\pi(2e^{-3\frac{1}{2}} + 1).$$

1.5 Show the area of A stability.

2 Conclusion

Thank you to Prof. Hamfeldt, neé Froese, for reading this work, for her instruction, lectures and future office hours efforts. I look forward to any feedback and learning more of the material in this course.

References

- [1] Kendall E Atkinson. *An introduction to numerical analysis*. John wiley & sons, 2008.

José Pabón