

Numerical Analysis Linear Algebra Previous Qual - August 2018.

José L. Pabón

Spring 2020

This work is based on the course textbook [1], the material discussed in lectures and office hours related to our course MAT614 and additional references.

1 Problem August 2018 1 -

1.1 1a

1.2 1b

2 Problem 2

2.1 2a

2.2 2b

3 Problem August 2018 3 - Solve this system of ODE IVP:

$$\dot{y} = 5y_1 - 6y_2.$$

$$\dot{y} = 3y_1 - 4y_2.$$

$$y_1(0) = 4, y_2(0) = 1$$

3.1 REDACTED

For IVP problems of this type, we have that our standard solutions relative to eigenvalues $\lambda = \mu + i\nu$ are:

$$y = e^{\mu t}(\alpha \sin(\nu t) + \beta \cos(\nu t)).$$

3.2 Solution proof

We have that our corresponding matrix representation for this system is:

$$A = \begin{bmatrix} 5 & -6 \\ 3 & -4 \end{bmatrix}$$

$$\begin{bmatrix} 5 & -6 \\ 3 & -4 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$

We calculate the eigenvalues of A by solving $\det(A - \lambda I) = 0 \implies (5 - \lambda)(-4 - \lambda) - (3(-6)) = 0 \implies \lambda^2 - \lambda - 2 = 0$. We use these two eigenvalues to solve $(A - \lambda I)v = 0$ to find the corresponding eigenvalues:

$$\text{For } \lambda_1 = 2, v_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

$$\text{For } \lambda_2 = -1, v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

We thus have our general solution:

$$y_{gen} = \alpha e^{2t} \begin{bmatrix} 2 \\ 1 \end{bmatrix} + \beta e^{-t} \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

We find our particular solution to this IVP by using the initial condition provided:

$$y_{part} = \alpha e^{20} \begin{bmatrix} 2 \\ 1 \end{bmatrix} + \beta e^{-0} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \end{bmatrix}.$$

We solve this system of equations to determine that $\alpha = 3, \beta = -2$.

Thus, our final particular solution to this particular initial value problem is:

$$y_{(part.final)} = 3e^{2t} \begin{bmatrix} 2 \\ 1 \end{bmatrix} - 2e^{-t} \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

Or, equivalently:

$$\therefore y_{(part.final)} = y_1 + y_2, y_1 = 6e^{2t} - 2e^{-t}, y_2 = 3e^{2t} - 2e^{-t}. \checkmark$$

4 Conclusion

Thank you to Prof. Hamfeldt, neé Froese, for reading this work, for her instruction, lectures and future office hours efforts. I look forward to any feedback and learning more of the material in this course.

References

- [1] Kendall E Atkinson. *An introduction to numerical analysis*. John wiley & sons, 2008.

José Pabón