

Numerical Analysis Previous Qual - August 2018.

José L. Pabón

Spring 2020

This work is based on the course textbook [1], the material discussed in lectures and office hours related to our course MAT614 and additional references.

1 Problem August 2018 6 - REDACTED

recision is less than or equal to $2n - 1$, and then that it is no more than the same.

Suppose inner product with weight is defined by:

$$\langle f, g \rangle = \int_a^b f(x)g(x)w(x)dx.$$

Consider quadrature formula:

$$I(f) = \int_a^b f(x)w(x)dx \approx Q(f) = \sum_{j=1}^n w_j f(x_j).$$

We have that:

$$w_j = \int_a^b l_j(x)w(x)dx.$$

And

$$l_j(x) = \prod_{i=1,2,3,\dots,n-i \neq j} \frac{(x - x_i)}{(x_j - x_i)}.$$

1.1 Solution, proof:

Degree of precision is defined as the highest order polynomial that the quadrature will return an exact answer for. Study note - for Simpsons rule this is three, for trapezoid rule this is one.

The key idea of this proof is division of the polynomial f . We define polynomials such that:

$$\frac{f(x)}{p_n(x)} - \frac{r(x)}{p_n(x)} = q(x) \implies f(x) = p_n(x)q(x) + r(x).$$

Where rational expression is shorthand for polynomial division. We construct this with the degrees of f, p, q, r being $\leq 2n - 1, n, n - 1, n - 1$ and by construction and the interpolation of $I(f)$, *★ we have that $r(x)$ is exact and there is no further remainder.*

★ \implies ASIDE note to the group - I went to Prof. Hamfeldt's office hour to talk about this problem, she mentioned this claim in italics is not clear or obvious. I asked her how I could formulate my argument better, she suggested 'expressing the remainder function $r(x)$ in terms of basis functions for the problem, and it should follow.' Still puzzling over this.

END ASIDE.

We have that $p_n x$ is orthogonal to all polynomials of degree $\leq n - 1$, thus we have that:

$$I(f) = I(qp_n + r) = \int_a^b q(x)p_n(x) + r(x)dx + \int_a^b r(x)w(x)dx = 0 + \int_a^b r(x)w(x)dx = Q(r).$$

Thus we find that:

$$I(f) = Q(r) = \sum w_j r(x_j) = \sum w_j (p_n(x_j)r(x_j) + r(x_j)) = \sum 0 + w_j r(x_j) = Q(f).$$

Need to show the remainder polynomials are exact.

Thus, we have that the precision of this method (Gaussian quadrature) is $\leq 2n - 1$.

1.2 Show that the precision is no greater than $2n - 1$.

We assume the same construction as in the previous argument, except for the degrees of the polynomials:

$$\frac{f(x)}{p_n(x)} - \frac{r(x)}{p_n(x)} = q(x) \implies f(x) = p_n(x)q(x) + r(x).$$

Where rational expression is shorthand for polynomial division. We construct this with the degrees of f, p, q, r being $\leq 2n, n, n, n - 1$ and by construction and the interpolation of $I(f)$, we have that $r(x)$ is still exact and there is no further remainder.

We have that $p_n x$ is orthogonal to all polynomials of degree $\leq n - 1$ but not orthogonal to all polynomials of degree n , thus we have that:

$$I(f) = I(qp_n + r) = \int_a^b q(x)p_n(x) + r(x)dx + \int_a^b r(x)w(x)dx = 0 + \int_a^b r(x)w(x)dx = Q(r) + \langle q, p_n \rangle.$$

Thus, $\langle q, p_n \rangle \neq 0$.

We still have the same zeros of our p_n , so thus we find that:

$$Q(f) = \sum w_j r(x_j) = Q(qp_n + r) \sum w_j (p_n(x_j)r(x_j) + r(x_j)) = \sum 0 + r(x_j) = Q(r).$$

Thus,

$$Q(f) \neq Q(r)$$

Thus, for all polynomials f of degree $2n$ or higher, the quadrature formula is not precise.

2 Conclusion

Thank you to Prof. Hamfeldt, neé Froese, for reading this work, for her instruction, lectures and future office hours efforts. I look forward to any feedback and learning more of the material in this course.

References

- [1] Kendall E Atkinson. *An introduction to numerical analysis*. John wiley & sons, 2008.