

CS 341, Fall 2013
Solutions for Midterm 1

1. (a) True. By HW 2, problem 3, we know \overline{A} is regular. Since \overline{A} and B are regular, then $\overline{A} \cup B$ is regular by Theorem 1.25. Theorem 1.49 then implies $(\overline{A} \cup B)^*$ is regular.
 - (b) False. See HW 6, problem 2(a).
 - (c) False. For example, $A = \{0^n 1^n 0^n \mid n \geq 0\}$ is a subset of $B = L((0 \cup 1)^*)$, but A is non-context-free and B is context-free.
 - (d) True. Use the pumping lemma with string $a^p b^p$, where $p \geq 3$ is the pumping length.
 - (e) True. Since A has a regular expression, A is a regular language by Theorem 1.54. Then Corollary 2.32 implies A is also context-free, so it has a CFG. Theorem 2.9 then ensures that A has a CFG in Chomsky normal form.
 - (f) True. Suppose A is non-context-free but regular. But then Corollary 2.32 implies A is context-free, which is a contradiction.
 - (g) True. See slide 2-111.
 - (h) True. HW 4, problem 5(a).
 - (i) True. HW 4, problem 5(c).
 - (j) False. 1^* is regular since it has a regular expression, but this language is infinite.
2. (a) $b^* a b^* \cup b^* a b^* (a \cup \varepsilon) b^*$. Another regular expression is $b^*(a \cup a b^*(a \cup \varepsilon)) b^*$. There are infinitely many regular expressions for the language.
 - (b) $G' = (V', \Sigma, R', S_0)$, where $V' = V \cup \{S_0\}$, S_0 is the (new) starting variable, Σ is the same alphabet of terminals as in G , and $R' = R \cup \{S_0 \rightarrow SS_0, S_0 \rightarrow \varepsilon\}$.
 - (c) $M_3 = (Q_3, \Sigma, \delta_3, q_3, F_3)$, where $Q_3 = Q_1 \times Q_2$; Σ is the same alphabet as M_1 and M_2 have; the transition function δ_3 satisfies $\delta((q, r), \ell) = (\delta_1(q, \ell), \delta_2(r, \ell))$ for $(q, r) \in Q_3$ and $\ell \in \Sigma$; the starting state $q_3 = (q_1, q_2)$; and $F_3 = (Q_1 \times F_2) \cap (F_1 \times Q_2)$, which also can be written as $F_1 \times F_2$.
 - (d) After one step, the CFG is then


$$\begin{aligned} S_0 &\rightarrow S \\ S &\rightarrow A1SA \mid 1SA \mid A1S \mid 1S \mid A0 \mid 0 \mid \varepsilon \\ A &\rightarrow 0S0 \end{aligned}$$

3. (a) A regular expression for L_1 is


$$R_1 = (+ \cup - \cup \varepsilon) \Sigma_1 \Sigma_1^*$$

where $\Sigma_1 = \{0, 1, 2, \dots, 9\}$ as defined in the problem.

(b) An NFA for L_1 is



(c) Define $\Sigma_2 = \{-, +\}$ and $\Sigma_3 = \Sigma_2 \cup \{.\}$. Then a DFA for L_1 is




(d) A regular expression for L_2 is

$$R_2 = (+ \cup - \cup \varepsilon)(\Sigma_1 \Sigma_1^* . \Sigma_1^* \cup . \Sigma_1 \Sigma_1^*)$$

Note that the regular expression $(+ \cup - \cup \varepsilon) \Sigma_1^* . \Sigma_1^*$ is not correct since it can generate the strings “.”, “+.” and “-.”, which are not valid floating-point numbers.


(e) An NFA for L_2 is



(f) Note that $L = L_1 \cup L_2$, so a regular expression for L is

$$R_3 = R_1 \cup R_2$$

(g) We can construct an NFA for L by taking the union of the NFA's for L_1 and L_2 as follows:




There are many other correct answers for this and the other parts.

4. (a) $G = (V, \Sigma, R, S)$ with set of variables $V = \{S, X\}$, where S is the start variable; set of terminals $\Sigma = \{a, b, c\}$; and rules

$$\begin{aligned} S &\rightarrow bSc \mid X \\ X &\rightarrow aXc \mid \varepsilon \end{aligned}$$

- (b) PDA



For every b and a read in the first part of the string, the PDA pushes an x onto the stack. Then it must read a c for each x popped off the stack.

5. Language A is nonregular. To prove this, suppose that A is a regular language. Let p be the “pumping length” of the Pumping Lemma. Consider the string $s = b^p a^{p+1}$. Note that $s \in A$, and $|s| = 2p + 1 \geq p$, so the Pumping Lemma will hold. Thus, there exists strings x , y , and z such that $s = xyz$ and

- (a) $xy^i z \in A$ for each $i \geq 0$,
- (b) $|y| > 0$,
- (c) $|xy| \leq p$.

Since the first p symbols of s are all b 's, the third property implies that x and y consist only of b 's. So z will be the rest of the b 's, followed by a^{p+1} . The second property states that $|y| > 0$, so y has at least one b . More precisely, we can then say that

$$\begin{aligned} x &= b^j \text{ for some } j \geq 0, \\ y &= b^k \text{ for some } k \geq 1, \\ z &= b^m a^{p+1} \text{ for some } m \geq 0. \end{aligned}$$

Since $b^p a^{p+1} = s = xyz = b^j b^k b^m a^{p+1} = b^{j+k+m} a^{p+1}$, we must have that $j + k + m = p$. The first property implies that $xy^2z \in A$, but

$$\begin{aligned} xy^2z &= b^j b^k b^k b^m a^{p+1} \\ &= b^{p+k} a^{p+1} \end{aligned}$$

since $j + k + m = p$. Hence, $xy^2z \notin A$ because it doesn't have more a 's than b 's since $k \geq 1$, and we get a contradiction. Therefore, A is a nonregular language.