

023 Recitation Notes F sept. 6 - Nick
1, 5, 8*, 9, 11, 13, 16

defs.


gen $V = \int_a^b A(x) dx$

disk $V = \int_a^b \pi r(x)^2 dx$

washer $V = \int_a^b \pi (R(x)^2 - r(x)^2) dx$

note $= \int_a^b \pi R(x)^2 dx - \int_a^b \pi r(x)^2 dx$



(1) solid lies between planes perpendicular to the x -axis at $x=0$ and $x=4$. Cross sections perpendicular to the x -axis on $x \in [0, 4]$ are squares whose diagonals run from the parabola $y = -\sqrt{x}$ and $y = \sqrt{x}$



(5) base of a solid

(a) equilateral triangle

is between the x -axis and the curve $y = 2\sqrt{\sin x}$ and $x \in [0, \pi]$



$$h^2 = 4\sin x - \sin x$$

$$h^2 = 3\sin x$$

$$h = \sqrt{3\sin x}$$

$$A(x) = \frac{1}{2}bh$$

$$= \frac{1}{2}(2\sqrt{\sin x})(\sqrt{3\sin x})$$

$$= \sqrt{3}\sin x$$

at x we have a square

$$a^2 + a^2 = 4x$$

$$2a^2 = (2\sqrt{x})^2$$

$$a = \frac{2\sqrt{x}}{\sqrt{2}}$$

$$a = \sqrt{2x}$$

$$A(x) = a^2$$


$$A(x) = 2x$$

$$V = \int_0^4 A(x) dx = \int_0^4 2x dx$$

$$V = \int_0^\pi A(x) dx =$$

$$V = \int_0^\pi \sqrt{3}\sin x dx$$



* (8)



base of a solid is in the region bounded by
 $y = \sqrt{x}$ and $y = \frac{x}{2}$

HW prob
just show
a little


(a) isosceles triangle of height 6



$$A(x) = \frac{1}{2} (\sqrt{x} - x/2)(6)$$

$$V = \int_0^2 3\sqrt{x} - \frac{3}{2}x \, dx$$

(b) semicircles whose diameters are the base



$$\begin{aligned} A(x) &= \frac{1}{2} \pi r^2 \\ &= \frac{1}{2} \pi \left(\frac{\sqrt{x} - x/2}{2}\right)^2 \\ &= \frac{1}{2} \cdot \frac{1}{4} \pi (x - x\sqrt{x} + \frac{x^2}{4}) \end{aligned}$$

$$V = \int_0^2 \frac{\pi}{8} (\frac{x^2}{4} - x\sqrt{x} + x) \, dx$$

$$= \frac{\pi}{8} \int_0^2 (\frac{x^2}{4} - x^{3/2} + x) \, dx$$

$$= \frac{\pi}{8} \left[\frac{1}{3} \frac{x^3}{4} - \frac{2}{5} x^{5/2} + \frac{1}{2} x^2 \right] \Big|_{x=0}^2$$


$$= \frac{\pi}{8} \left[\frac{1}{12}(8) - \frac{2}{5}(4\sqrt{2}) + \frac{(4)}{2} \right] - [0 - 0 + 0]$$

$$= \pi \left[\frac{1}{12} - \frac{(2)(4)}{8} \left(\frac{4\sqrt{2}}{5} \right) + \frac{2}{8} \right]$$

$$= \pi \left(\frac{1}{3} - \frac{\sqrt{2}}{5} \right)$$

②

(a) Solid lines between planes perpendicular to the y -axis $y \in [0, 2]$ are circular disks with diameter running from the y -axis to the parabola $x = \sqrt{5}y^2$




$$A(y) = \pi \left(\frac{\sqrt{5}}{2} y^2 \right)^2$$

$$V = \int_0^2 \pi \left(\frac{\sqrt{5}}{2} y^2 \right)^2 dy$$


$$= \int_0^2 \frac{\pi}{4} \cdot 5 y^4 dy$$

(II) right-tetrahedron volume



$$V = \int_0^5 A(x) dx$$

$$\begin{aligned} A(x) &= \frac{1}{2} b h \\ &= \frac{1}{2} b(x) h(x) \end{aligned}$$



③

(ii) cont...

$$\begin{array}{l} b(0) = 0 \\ b(5) = 4 \end{array} \quad \left. \begin{array}{l} \text{linear and passes thru. the origin} \\ b(x) = \frac{4}{5}x \end{array} \right.$$
$$\begin{array}{l} h(0) = 0 \\ h(5) = 3 \end{array} \quad \left. \begin{array}{l} h(x) = \frac{3}{5}x \end{array} \right.$$

$$A(x) = \frac{1}{2} \left(\frac{4}{5}x \right) \left(\frac{3}{5}x \right)$$
$$= \frac{12}{50} x^2 = \frac{6}{25} x^2$$

$$F = \int_0^{25} \frac{12}{50} x^2 dx$$

$$= \left(\frac{6}{25} \right) \left[\frac{1}{3} x^3 \right] \Big|_{x=0}^{25}$$



$$= \left(\frac{6}{25} \right) \left(\frac{1}{3} \right) (125 - 0)$$

$$F = 2$$

④

(13) twisted solid - square of side length s lies in a plane perpendicular to a line L and one of the vertices touches the line. Square sweeps out a distance h along L and simultaneously completes a full turn about the line

(a) the volume of the swept solid
 (b) what if it makes two full turns



$$A(x) = s^2$$


$$V = \int_0^h A(x) dx$$

$$= \int_0^h s^2 dx$$


$$V = s^2 h$$

what about (b)


(16) Gasoline in a tank



(a)



circle area




$$\frac{1}{2} = \frac{180}{360} \quad \frac{1}{4} = \frac{90}{360}$$

$$A_1 = \frac{\theta}{360} \pi r^2$$

$$A_2 = \frac{1}{2} b h$$

$$r = 4 \text{ ft}$$

$$h = 2 \text{ ft}$$



$$\alpha = \frac{1}{2} \alpha$$

$$\theta = 360 - \alpha$$

Soh Cah Toa

$$\frac{2}{4} = \cos(\alpha)$$

$$\frac{1}{2} = \cos(\alpha)$$

$$\cos(1/2) = 60^\circ$$

triangle area

$$\alpha = 120^\circ \quad \theta = 240^\circ$$





$$b = 2c$$

$$\frac{c}{4} = \sin(\alpha) = \sqrt{3}/3$$

$$b = 2(4\sqrt{3}/3)$$

(6)

(16) cont... (b)



$$A(y) = 2(10\text{ft}) z(y)$$

$$z(y) = ?$$

$$y^2 + z(y)^2 = (4\text{ft})^2$$

$$z(y)^2 = (4\text{ft})^2 - y^2$$

$$z(y) = \sqrt{16\text{ft}^2 - y^2}$$

$$A(y) = (20\text{ft}) \sqrt{(16\text{ft}^2) - y^2}$$

$$T = \int_{-4}^2 (20\text{ft}) \sqrt{(16\text{ft}^2) - y^2} dy$$

7

