

Web-based Supplementary Materials for
“Procedures Controlling the k -FDR Using Bivariate Distributions
of the Null p -Values”

Sanat K. Sarkar and Wenge Guo

Temple University and National Institute of Environmental Health Sciences

Web Appendix A: Proofs

PROOF OF LEMMA 2.1. Let $\hat{P}_{(1)} \leq \dots \leq \hat{P}_{(n_0)}$ be the ordered values of $\{\hat{P}_1, \dots, \hat{P}_{n_0}\}$. Note that $\hat{P}_{(j)} \leq P_{(n-n_0+j)}$ for all $1 \leq j \leq n_0$. Therefore, the original stepwise procedure rejects less number of null hypotheses, and hence has less number of falsely rejected null hypotheses, than the corresponding stepwise procedure where the p -values corresponding to the false null hypotheses are all very close to zero. Thus, the lemma follows. ■

PROOF OF THEOREM 3.1.

$$\begin{aligned}
 k\text{-FDR} &= E \left\{ \frac{V}{R} \cdot I(V \geq k) \right\} = E \left\{ \sum_{r=k}^n \frac{1}{r} \sum_{i=1}^{n_0} I(\hat{P}_i \leq \alpha_r, V \geq k, R = r) \right\} \\
 &= \sum_{i=1}^{n_0} \sum_{r=k}^n \frac{1}{r} \Pr \left\{ \hat{P}_i \leq \alpha_r, V \geq k, R = r \right\} \\
 &= \frac{\alpha_k}{k} \sum_{i=1}^{n_0} \sum_{r=k}^n P \left\{ V \geq k, R = r \mid \hat{P}_i \leq \alpha_r \right\}. \tag{A.1}
 \end{aligned}$$

Now, for any $r > k$ and $i \leq n_0$,

$$\begin{aligned}
 &\Pr \left\{ V \geq k, R = r \mid \hat{P}_i \leq \alpha_r \right\} \\
 &= \Pr \left\{ V \geq k, R \geq r \mid \hat{P}_i \leq \alpha_r \right\} - \Pr \left\{ V \geq k, R \geq r+1 \mid \hat{P}_i \leq \alpha_r \right\} \\
 &\leq \Pr \left\{ V \geq k, R \geq r \mid \hat{P}_i \leq \alpha_{r-1} \right\} - \Pr \left\{ V \geq k, R \geq r+1 \mid \hat{P}_i \leq \alpha_r \right\}.
 \end{aligned}$$

The inequality follows from the assumption (1), since $\{V \geq k, R \geq r\}$ is a decreasing set for a

stepwise procedure. Thus, we get

$$\begin{aligned}
k\text{-FDR} &\leq \frac{\alpha_k}{k} \sum_{i=1}^{n_0} \Pr \left\{ V \geq k, R = k \mid \hat{P}_i \leq \alpha_k \right\} + \\
&\quad \frac{\alpha_k}{k} \sum_{i=1}^{n_0} \sum_{r=k+1}^n \Pr \left\{ V \geq k, R = r \mid \hat{P}_i \leq \alpha_r \right\} \\
&\leq \frac{\alpha_k}{k} \sum_{i=1}^{n_0} \Pr \left\{ V \geq k, R = k \mid \hat{P}_i \leq \alpha_k \right\} + \\
&\quad \frac{\alpha_k}{k} \sum_{i=1}^{n_0} \Pr \left\{ V \geq k, R \geq k+1 \mid \hat{P}_i \leq \alpha_k \right\} \\
&= \frac{\alpha_k}{k} \sum_{i=1}^{n_0} \Pr \left\{ V \geq k \mid \hat{P}_i \leq \alpha_k \right\} = \frac{1}{k} \sum_{i=1}^{n_0} \Pr \left\{ V \geq k, \hat{P}_i \leq \alpha_k \right\}. \tag{A.2}
\end{aligned}$$

Applying Lemma 2.1 to (A.2), we get

$$k\text{-FDR} \leq \frac{1}{k} \sum_{i=1}^{n_0} P \left\{ \hat{R}_{n_0} \geq k, \hat{P}_i \leq \alpha_k \right\}.$$

Let $\hat{R}_{n_0-1}^{(-i)}$ denote the number of rejections in the corresponding stepwise procedure based on the ordered values $\hat{P}_{(1)}^{(-i)} \leq \dots \leq \hat{P}_{(n_0-1)}^{(-i)}$ of $\{\hat{P}_1, \dots, \hat{P}_{n_0}\} \setminus \{\hat{P}_i\}$ and the n_0-1 critical values $\alpha_{n-n_0+2} \leq \dots \leq \alpha_n$. Then, for any $k \leq r \leq n_0$, we notice that, when our procedure is a stepup procedure

$$\begin{aligned}
&\left\{ \hat{R}_{n_0} = r, \hat{P}_i \leq \alpha_k \right\} \\
&= \left\{ \hat{P}_{(r)} \leq \alpha_{n-n_0+r}, \hat{P}_{(r+1)} > \alpha_{n-n_0+r+1}, \dots, \hat{P}_{(n_0)} > \alpha_n, \hat{P}_i \leq \alpha_k \right\} \\
&= \left\{ \hat{P}_{(r-1)}^{(-i)} \leq \alpha_{n-n_0+r}, \hat{P}_{(r)}^{(-i)} > \alpha_{n-n_0+r+1}, \dots, \hat{P}_{(n_0-1)}^{(-i)} > \alpha_n, \hat{P}_i \leq \alpha_k \right\} \\
&= \left\{ \hat{R}_{n_0-1}^{(-i)} = r-1, \hat{P}_i \leq \alpha_k \right\}; \tag{A.3}
\end{aligned}$$

whereas, for a stepdown procedure, we have

$$\begin{aligned}
& \left\{ \hat{R}_{n_0} = r, \hat{P}_i \leq \alpha_k \right\} \\
&= \left\{ \hat{P}_{(1)} \leq \alpha_{n-n_0+1}, \dots, \hat{P}_{(r)} \leq \alpha_{n-n_0+r}, \hat{P}_{(r+1)} > \alpha_{n-n_0+r+1}, \hat{P}_i \leq \alpha_k \right\} \\
&\subseteq \left\{ \hat{P}_{(1)}^{(-i)} \leq \alpha_{n-n_0+2}, \dots, \hat{P}_{(r-1)}^{(-i)} \leq \alpha_{n-n_0+r}, \hat{P}_{(r)}^{(-i)} > \alpha_{n-n_0+r+1}, \hat{P}_i \leq \alpha_k \right\} \\
&= \left\{ \hat{R}_{n_0-1}^{(-i)} = r-1, \hat{P}_i \leq \alpha_k \right\}. \tag{A.4}
\end{aligned}$$

Thus, for a stepwise (stepup or stepdown) procedure, we get

$$\begin{aligned}
k\text{-FDR} &\leq \frac{1}{k} \sum_{i=1}^{n_0} \sum_{r=k}^{n_0} \Pr \left\{ \hat{R}_{n_0-1}^{(-i)} = r-1, \hat{P}_i \leq \alpha_k \right\} \\
&= \frac{1}{k} \sum_{i=1}^{n_0} \sum_{j(\neq i)=1}^{n_0} \sum_{r=k}^{n_0} \frac{1}{r-1} \Pr \left\{ \hat{R}_{n_0-1}^{(-i)} = r-1, \hat{P}_i \leq \alpha_k, \hat{P}_j \leq \alpha_{n-n_0+r} \right\}. \tag{A.5}
\end{aligned}$$

As explained in (A.3) and (A.4),

$$\left\{ \hat{R}_{n_0-1}^{(-i)} = r-1, \hat{P}_i \leq \alpha_k, \hat{P}_j \leq \alpha_{n-n_0+r} \right\} = \left\{ \hat{R}_{n_0-2}^{(-i,-j)} = r-2, \hat{P}_i \leq \alpha_k, \hat{P}_j \leq \alpha_{n-n_0+r} \right\},$$

for a stepup procedure, and

$$\left\{ \hat{R}_{n_0-1}^{(-i)} = r-1, \hat{P}_i \leq \alpha_k, \hat{P}_j \leq \alpha_{n-n_0+r} \right\} \subseteq \left\{ \hat{R}_{n_0-2}^{(-i,-j)} = r-2, \hat{P}_i \leq \alpha_k, \hat{P}_j \leq \alpha_{n-n_0+r} \right\},$$

for a stepdown procedure, where $\hat{R}_{n_0-2}^{(-i,-j)}$ is the number of rejections in the corresponding stepwise procedure involving $\left\{ \hat{P}_1, \dots, \hat{P}_{n_0} \right\} \setminus \left\{ \hat{P}_i, \hat{P}_j \right\}$ and critical values $\alpha_{n-n_0+3} \leq \dots \leq \alpha_n$. Thus,

$$k\text{-FDR} \leq \frac{(n-n_0+k)\alpha_k}{k^2(k-1)} \sum_{i=1}^{n_0} \sum_{j(\neq i)=1}^{n_0} \sum_{r=k}^{n_0} \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} = r-2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r} \right\}.$$

The inequality follows from the fact that

$$\frac{n - n_0 + r}{r - 1} \leq \frac{n - n_0 + k}{k - 1},$$

for all $k \leq r \leq n_0$. Since for $r > k$,

$$\begin{aligned} & \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} = r - 2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r} \right\} \\ = & \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} \geq r - 2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r} \right\} - \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} \geq r - 1, \right. \\ & \left. \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r} \right\} \\ \leq & \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} \geq r - 2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r-1} \right\} - \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} \geq r - 1, \right. \\ & \left. \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r} \right\}, \end{aligned}$$

with the inequality following due to the property (1) and the fact that $\{\hat{R}_{n_0-2}^{(-i,-j)} \geq r - 2, \hat{P}_i \leq \alpha_k\}$

is decreasing, we have

$$\begin{aligned} & \sum_{r=k}^{n_0} \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} = r - 2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r} \right\} \\ \leq & \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} = k - 2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+k} \right\} + \\ & \sum_{r=k+1}^{n_0} \left[\Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} \geq r - 2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r-1} \right\} - \right. \\ & \left. \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} \geq r - 1, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+r} \right\} \right] \\ = & \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} = k - 2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+k} \right\} + \\ & \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} \geq k - 1, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+k} \right\} \\ = & \Pr \left\{ \hat{R}_{n_0-2}^{(-i,-j)} \geq k - 2, \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+k} \right\}. \end{aligned}$$

Therefore,

$$\begin{aligned}
& k\text{-FDR} \\
& \leq \frac{(n - n_0 + k)\alpha_k}{k^2(k - 1)} \sum_{i=1}^{n_0} \sum_{j(\neq i)=1}^{n_0} \Pr \left\{ \hat{R}_{n_0-2}^{(-i, -j)} \geq k - 2, \hat{P}_i \leq \alpha_k \mid \right. \\
& \quad \left. \hat{P}_j \leq \alpha_{n-n_0+k} \right\} \\
& \leq \frac{(n - n_0 + k)\alpha_k}{k^2(k - 1)} \sum_{i=1}^{n_0} \sum_{j(\neq i)=1}^{n_0} \Pr \left\{ \hat{P}_i \leq \alpha_k \mid \hat{P}_j \leq \alpha_{n-n_0+k} \right\} \\
& = \frac{1}{k(k - 1)} \sum_{i=1}^{n_0} \sum_{j(\neq i)=1}^{n_0} \Pr \left\{ \hat{P}_i \leq \alpha_k, \hat{P}_j \leq \alpha_{n-n_0+k} \right\}. \tag{A.6}
\end{aligned}$$

The theorem then follows by considering the maximum of the right-hand side in (A.6) over the set of values of n_0 . ■

PROOF OF (8). Since the p -values are independent and $\alpha_i = \{i \vee k\}\beta/n$, $i = 1, \dots, n$, we see from the first line in (A.5) that

$$k\text{-FDR} \leq \frac{n_0\beta}{n} \sum_{r=k}^{n_0} \Pr \left\{ \hat{R}_{n_0-1} = r - 1 \right\} = \frac{n_0\beta}{n} \Pr \left\{ \hat{R}_{n_0-1} \geq k - 1 \right\},$$

the desired inequality. ■

PROOF OF THEOREM 5.1. Define $p_{ijr} = P \left\{ \hat{P}_i \in [\alpha_{j-1}, \alpha_j], V \geq k, R = r \right\}$, given a set critical

values $0 = \alpha_0 < \alpha_1 < \dots < \alpha_n$. Then, from (A.1) we have,

$$\begin{aligned}
k\text{-FDR} &= \sum_{r=k}^n \sum_{i=1}^{n_0} \frac{1}{r} \Pr \left\{ \hat{P}_i \leq \alpha_r, V \geq k, R = r \right\} = \sum_{r=k}^n \sum_{i=1}^{n_0} \sum_{j=1}^r \frac{1}{r} p_{ijr} \\
&= \sum_{i=1}^{n_0} \sum_{j=1}^n \sum_{r=k \vee j}^n \frac{1}{r} p_{ijr} \leq \sum_{i=1}^{n_0} \sum_{j=1}^n \frac{1}{k \vee j} \sum_{r=k \vee j}^n p_{ijr} \\
&\leq \sum_{i=1}^{n_0} \sum_{j=1}^n \frac{1}{k \vee j} \Pr \left\{ \hat{P}_i \in [\alpha_{j-1}, \alpha_j], V \geq k \right\} \\
&\leq \sum_{i=1}^{n_0} \sum_{j=1}^n \frac{\alpha_j - \alpha_{j-1}}{k \vee j}.
\end{aligned}$$

Let $\alpha_i = (i \vee k) \alpha_k / k$, for some fixed $0 < \alpha_k < 1$. Then,

$$k\text{-FDR} \leq n_0 \frac{\alpha_k}{k} \left\{ 1 + \sum_{j=k+1}^n \frac{1}{j} \right\}. \quad (\text{A.7})$$

The theorem then follows by considering $\alpha_k = k\alpha/n \left\{ 1 + \sum_{j=k+1}^n \frac{1}{j} \right\}$ in (A.7).

■

Web Figure 1:




Figure 1: Power of two k -FDR stepdown procedures in the case of independence with parameters $n = 200, k = 5, d = 2$ and $\alpha = 0.05$.